Monmouth Gastroenterology, LLC

A Division of Allied Digestive Health 1912 Hwy 35 South, Suite 201 Oakhurst, NJ 07755 732-389-5004 fax 732-389-1850

Doctors: Baig, Belitsis, Fiest, Gorcey, Merikhi and Uppal

Date:	Time: THE FACILITY WILL CALL YOU WITH THE EXACT TIME EITHER ONE
OR TWO DAYS PRIOR	TO YOUR PROCEDURE.
Required Preparation for Your Colonoscopy – MOVIPREP-SPLIT: 11AM TO 1:50PM Please follow the below instructions—not the instructions printed on the package	
and did not tell the doctor, Vitamin E, Fish Oil, Garlic T etc.) for 5 days prior to the Xarelto, Pradaxa, Eliquis a HEADACHE OR PAIN.	cheduled Colonoscopy: If you have diabetes or take blood thinners (Coumadin, Plavix, Ticlid, Aspirin, etc.) inform the doctor immediately. Generally you should stop medications, or vitamins containing Iron, ablets, Ginko Biloba, ginseng or Aspirin-like products (Advil, Motrin, Aleve, Nuprin, Alka-Seltzer, Naprosyn, procedure. However, you will be specifically instructed about withholding Aspirin, Plavix, Warfarin, and Coumadin according to your medical situation. YOU MAY TAKE TYLENOL AS NEEDED FOR A TICPLEASE MAKE SCHEDULER AWARE – IN ORDER TO RECEIVE INSTRUCTIONS ON YOUR MEDICATION/ INSULIN
<pre>procedure). This is essent</pre>	nas and melon are permitted), nuts, vegetables, salads, seeds, berries or whole grains (until after the ial in order to successfully clear your colon for the procedure.
 Have a regular brea From noon until 6 projuice, Gatorade, lend INGEST NOTHING At 8:00pm mix firstop of the contained reconstituted solution At 8:30 PM Drink firstop of the contained reconstituted solution At 8:30 PM Drink firstop of the contained reconstituted solution At 5:30 AM mixtop of the contained reconstituted solution At 6:00 AM Drink firstop of the contained reconstituted solution At 6:00 AM Drink firstop of the contained reconstituted solution At 6:00 AM Drink firstop of the contained reconstituted solution 	second dose, empty 1 Pouch A and 1 Pouch B into the disposable container. Add lukewarm drinking water einer. Mix to dissolve. <i>If preferred, mix solution ahead of time and refrigerate prior to drinking. The should be used within 24 hours.</i> k second dose. The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to ely 8 oz), until the full liter is consumed. Drink 16 oz of the clear liquid of your choice. This is a necessary atte hydration and effective prep. MUST FINISH BY 7:00 AM.
	E: YOU NEED TO FINISH DRINKING YOUR LIQUID PREP 4 HOURS PRIOR TO YOUR TAKE YOUR HEART MEDICATIONS OR BLOOD PRESSURE MEDICATIONS
Ao	Please report to the following location: dvanced Endoscopy and Surgical Center: 142 Rte. 35, Eatontown, NJ732-935-0031

Monmouth Medical Center: 300 Second Avenue, Long Branch,NJ- Arrival time: _____outpatient registration Jersey Shore University Medical Center: 1945 Rte. 33, Neptune,NJ-Arrival time: ____outpatient registration