

Monmouth Gastroenterology, LLC
A Division of Allied Digestive Health 1912 Hwy 35 South, Suite 201 Oakhurst, NJ 07755 732-389-5004 fax 732-389-1850

Doctors: Baig, Belitsis, Fiest, Gorcey, Merikhi and Uppal

PATIEN	T NAME:Date:	
Time:	THE FACILITY WILL CALL YOU WITH THE EXACT TIME EITHER ONE OR TWO DAYS PRIOR TO YOUR	
	DURE.	
	NULYTELY/TRILYTE/COLYTE - PREPARATION FOR COLONOSCOPY—AM OR PM	
	Please follow the below instructions—not the instructions printed on the package	
FIVE DA	YS BEFORE Your Scheduled Colonoscopy: If you have diabetes or take blood thinners (Coumadin, Plavix, Ticlid, Aspirin, et	<b>c</b> 1
and did	not tell the doctor, inform the doctor immediately. Generally you should stop medications, or vitamins containing <b>Iron</b> , E, Fish Oil, Garlic Tablets, Ginko Biloba, ginseng or Aspirin-like products (Advil, Motrin, Aleve, Nuprin, Alka-Seltzer, Napros	
	5 days prior to the procedure. However, you will be specifically instructed about withholding Aspirin, Plavix, Warfarin,	•
Xarelto	Pradaxa, Eliquis and Coumadin according to your medical situation. YOU MAY TAKE TYLENOL AS NEEDED FOR A	
	CHE OR PAIN.	
IF Y	OU ARE A DIABETICPLEASE MAKE SCHEDULER AWARE – IN ORDER TO RECEIVE INSTRUCTIONS ON YOUR MEDICATION/ INSULIN	
TWO DA	YS BEFORE THE COLONOSCOPY: Do not eat any fruits (bananas and melon are permitted), nuts, vegetables, salads, seeds, berries or	
	ains (until after the procedure). This is essential in order to successfully clear your colon for the procedure.	
	/ BEFORE THE COLONOSCOPY:	
1. 2.	Mix PREP with water early in the day. Shake the container well and put it in the refrigerator.  Have a light breakfast and lunch – but finish eating by 12 noon. <b>DO NOT EAT DINNER!</b>	
2. 3.	From noon until 6:00 pm you may have <u>CLEAR LIQUIDS ONLY-You may have plain pasta</u> . Clear liquids include: tea, soft drinks, broth,	
Э.	juice (apple/ white grape), jello (avoid cherry /orange), AVOID MILK AND SOUPS.	
	DO NOT HAVE ANYTHING RED OR ORANGE IN COLOR (Colors effect what the colonoscopy sees.)	
4.	If ONLY stated by doctor –take 2 dulcolax tablets with 8ozs of water –check yes	
5.	YOU MAY CONTINUE CLEAR LIQUIDS UNTIL MIDNIGHT.	
DIRECTIO	ONS FOR COLONOSCOPY – SPLIT PREP –BEFORE 1:00 P.M.	
	-At 6:00 PM the night before the procedure, start to drink the liquid PREP. Drink 1/3 cup every five minutes until you finish drinking 2	
	liters. Place the other 2 liters into the fridge until you are ready to drink the next batch.	
В.	Start to drink the other 2 liters drink at (6 HOURS PRIOR TO YOUR PROCEDURE TIME). Drink 1/3 cup every five minutes unt	I
	you finish drinking the last 2 liters. You must finish drinking this prep 4 hours prior to your test.	
	owel movements are not clear (clear means yellow-greenish waterish movement with no solid stool or pieces) after you have finished t	ıe
	nen take two (2) Dulcolax tablets with 8 oz of clear liquid. You may have small amounts of clear liquids until midnight.  DNS FOR COLONOSCOPY- SAME DAY PREP-AFTER 1:00 P.M.	
1.	Day of the procedure – DO NOT EAT OR DRINK ANYTHING OTHER THAN YOUR PREP AND MEDICATIONS APPROVED BY YOUR DOCTO	R.
	-At(6 HOURS PRIOR TO YOUR PROCEDURE TIME) start to drink the liquid PREP – drink 1/3 cup every five minutesIf you	
	have completed most of the gallon PREP and have not had a bowel movement, you may use a fleet enema to induce a bowel movement	
	<u>Day of Procedure:</u> YOU NEED TO FINISH DRINKING YOUR LIQUID PREP 4 HOURS PRIOR TO YOUR PROCEDUR	Е <u>.</u>
	TAKE YOUR HEART MEDICATIONS OR BLOOD PRESSURE MEDICATIONS	
	Diago report to the following legation:	
	Please report to the following location:  Advanced Endoscopy and Surgical Center: 142 Rte. 35, Eatontown, NJ732-935-0031	
	Monmouth Medical Center: 300 Second Avenue, Long Branch, NJ- Arrival time: outpatient registratio	•
	Jersey Shore University Medical Center: 1945 Rte. 33, Neptune, NJ-Arrival time:outpatient registratio	